



Supplementary Materials for

Machine learning predicts which rivers, streams, and wetlands the Clean Water Act regulates

Simon Greenhill *et al.*

Corresponding authors: Simon Greenhill, sgreenhill@berkeley.edu; Joseph S. Shapiro, joseph.shapiro@berkeley.edu

Science **383**, 406 (2024)
DOI: [10.1126/science.adi3794](https://doi.org/10.1126/science.adi3794)

The PDF file includes:

Materials and Methods
Supplementary Text
Figs. S1 to S10
Tables S1 to S10
References

Other Supplementary Material for this manuscript includes the following:

MDAR Reproducibility Checklist

A. Materials and Methods

A.1 Model Choice, Design, and Assessment

5 We use ResNet-18 for a few reasons. Given the sample size for training data, additional blocks in the model may not improve performance. We observed some signs of overfitting in training data for ResNet-18, suggesting that the model is sufficiently complex and implying that a more complex model could exacerbate overfitting. We experimented with multi-task learning across binary regulation and the nine hydrological and the nine legal types of waters (SM section A.4) but found it did not improve accuracy in the validation data.

10 We also considered alternatives to ResNet but concluded they were unlikely to improve model performance. To improve model accuracy, meta-learning models may require substantially more data than we have. A foundation model would need to be pre-trained on large amounts of pre-existing data from a similar problem. In our setting, such data are not readily available. We experimented with a form of convolutional kernel ridge regression (37), a gradient boosting algorithm, and separate ResNet models for each rule, but found they did not have better validation accuracy than the model the main text uses.

15 Permutation importance tests, discussed in the main text, have limitations. A feature that is highly correlated with others may appear unimportant when removed, since the model can rely on the unpermuted features to arrive at the same prediction. We try to mitigate this by permuting features in groups, such as all PRISM or all Gridded National Soil Survey Geographic Database (gNATSGO) layers. Permutation can also produce unrealistic model inputs that generate unpredictable model behavior. Combined with the case studies and maps, however, these tests can clarify WOTUS-ML's functioning.

A.2 Train-Test Split

20 To prevent spatial leakage, we group data so AJDs in the same project or with geographically overlapping footprints are in the same fold. AJDs in the same project are evaluated by the same ACE engineer at the same time, so have correlated outcomes. Grouping by aerial imagery footprint ensures the test set does not partially include images and other input layers used to train the model.

25 We seek to assign an 80/10/10 split between train, test, and validation, stratifying the randomization by district-rule to ensure the same split in every district and every rule. Because groups span rules and districts, however, and we have limited data for some district-rules, this ideal split is not feasible in our setting.

30 We therefore adopt a splitting procedure that ensures we have sufficient data for testing and validation in all district-rules while getting as close as possible to the 80/10/10 target. We assign all the data from district-rules with 50 or fewer observations to test. District-rules with 100 or fewer observations are split 80/20 between test and validation. For district-rules with 500 or fewer observations, we assign 50 observations to test, then split the remaining observations 80-20 between training and validation. After the small district-rules are assigned, we split the remaining groups 80/10/10 between training, validation, and testing. We repeat the randomization until every district-rule with over 500 observations has at least 100 training

45 observations, and at least 5% of the observations in the validation and testing sets. The entire
randomization is stratified at the district level.¹

The maximum number of points in a single image footprint group is 244. The average is 3.67,
with a standard deviation of 8.53.

50

A.3 Input data

WOTUS-ML takes as inputs 34 layers from 11 distinct datasets (Table S3). These include
imagery from the National Agriculture Imagery Program (NAIP); geographic data on the
55 locations and characteristics of wetlands and streams from the National Wetlands Inventory
(NWI) and National Hydrography Dataset (NHD); soil characteristics from the Gridded National
Soil Survey Geographic Database; information on local climate from the PRISM Climate
Group's 30-year Normals; land cover from the National Land Cover Database (NLCD);
elevation from USGS's 3DEP model; ecoregion classification (level IV) from the US EPA;
60 distance to U.S. Army Corps of Engineers (ACE) district headquarters; and encodings for states,
ACE regulatory districts, and WOTUS rules. Several input layers are available only in the
contiguous US (CONUS), thus restricting our analysis to this region.

While state and ACE district layers represent political rather than geophysical inputs, they have
65 an important influence on what the CWA regulates. For example, if the St. Paul ACE district
concludes that no wetlands are regulated, and the Wilmington ACE district concludes that all
wetlands are regulated, the resulting AJDs have legally binding force and describe what the
CWA regulates. Similarly, we include a layer for the distance of a site to the closest ACE
70 headquarters because the probability of in-person visits decreases with this distance, and sites
with in-person visits have higher probability of being regulated. One potential explanation is that
field visits are more likely for sites nearer to an ACE office, and field visits may discover water
resources (e.g., swales under tree cover) that a desk analysis is less likely to observe. Although
CWA represents federal regulation, its management through ACE district offices implies that its
75 implementation is to some extent sub-national.

75

We rasterize all vector layers into 512x512 pixel tiles, where each pixel is 0.6 to 1.1 meters, to
match the resolution of the NAIP imagery. For the input layers that are not natively in a raster or
vector format (i.e., distance to ACE district headquarters and encodings for ACE districts, states,
and WOTUS rules), we fill a 512x512 pixel raster with one value to make these layers conform
80 to our convolutional neural network (CNN) architecture. For categorical data, such as the NLCD
land cover classes or ACE districts, we use ordinal encoding. We assign numeric values such
that similar classes or adjacent geographic areas are close together in numeric space (e.g., for
states, we do not use federal information processing standards (FIPS) codes, but instead assign
sequential numbers to neighboring states). After experimenting with one-hot encoding and
85 observing that it had no impact on accuracy in the validation set, we elect to use ordinal encoding
to avoid inputting an excessive number of layers into the CNN given the limited amount of

¹ Groups of observations occasionally overlap districts. A total of 61 out of the 14,000 image footprints groups span more than one district, accounting for about 6% of our observations. In cases where a group is assigned to multiple splits, we re-assign it to a single split, with train taking precedence over test and validation and test taking precedence over validation.

training data (~150,000 observations) and computational cost of training a model with more input layers. We use one-hot encoding for the rules, so each rule has its own input layer.

90 For the training data layers that contain information on built-up land (NAIP and NLCD), we use
images from the most recent survey *prior* to each AJD to ensure that the model's prediction is
based on pre-AJD conditions rather than post-AJD development activity. We use the most recent
survey available for the input layers corresponding to the four million prediction points across
95 the country for which we predict regulatory probabilities. We exclude all AJD and prediction
points that are missing input data, such as NAIP imagery or state information. All these excluded
AJDs are in the oceans, Great Lakes, or military bases (e.g., Area 51).

100 We highlight a few other notes on NHD and NWI. NHD describes 3.15 million stream miles in
the CONUS. We define a sample point as in NHD or NWI if it is within 10 meters of those
resources. Documentation for NHD's high resolution version states that it distinguishes ephemeral
from intermittent streams. We do not use the high-resolution NHD files given its incompleteness
and measurement error. NHD high resolution's distinction between ephemeral and intermittent
streams appears in only a few western states; elsewhere, NHD classifies all streams as intermittent.
Additionally, we found substantial differences between NHD's classification of streams as
intermittent or ephemeral and AJDs' classification of the same streams as intermittent or
105 ephemeral.

NWI and NHD seek to identify water types of locations across the US. While this has similar
outputs as the model, NWI and NHD alone provide insufficient information to determine whether
a site is jurisdictional (7). For example, NWI identifies wetlands, but does not identify whether a
wetland is abutting navigable waters or isolated, and thus does not identify if the wetlands are
110 jurisdictional. Similarly, while the high resolution NHD identifies certain water resources as
ephemeral (which would not be jurisdictional under NWPR), ACE may determine that some of
those water resources are in fact intermittent and thus jurisdictional under NWPR. Additionally,
NWI and NHD have imperfect measures, particularly for certain water types.

115 These considerations lead us to use NWI and NHD as model inputs, since they provide useful but
imperfect and incomplete information about regulatory probabilities. They may also affect the
model's accuracy. For example, if some future version of NHD and NWI had no errors, then those
error-free inputs would provide more useful information for a model, and thus a model trained on
perfect versions of NHD and NWI could have greater accuracy than WOTUS-ML.

120 At the same time, these considerations do not bear on our conclusions about model accuracy,
reliability, or bias. Our estimates of accuracy and bias are conditional on the information available
in the model inputs. Although using an NWI or NHD point alone may provide a biased estimate
of regulatory probabilities, by using NWI, NHD, and many other inputs to flexibly predict legally-
binding AJDs, the model obtains unbiased estimates of regulatory probabilities (Fig 2A).

125 Because PJDs are not certain to be jurisdictional, we use the model to assess their probability of
regulation, but do not train the model on PJDs. In other words, our data on PJDs do not affect
model performance since the model is trained only on AJDs. After training WOTUS-ML, we
compare predicted regulatory probabilities for PJDs versus other types of points. Sites with the
highest probability of regulation may be more likely to request PJDs than AJDs, so AJDs may

130 over-represent cases with ambiguous jurisdiction. We obtained details on PJDs from a Freedom of Information Act request filed with the EPA.

A.4 Other data

135 A few additional details on AJDs are useful to highlight. The ground-truth data include AJDs between August 2015 and May 2022, obtained from the EPA (38, 39). Of these, 81,316 were made under *Rapanos*, 61,582 under NWPR, and 7,755 under CWR. *Rapanos* applied through 2018; CWR applied through late 2020, though only in about half of states due to litigation; NWPR applied through mid-2021; and *Rapanos* applied thereafter. For a small subset of AJDs, we can manually obtain a PDF document for each AJD with modest additional detail on other variables from ACE's ORM2 database. A further subset of these PDF files includes idiosyncratic information from ACE engineers on what information the engineer gathered.

140 The main text mentions that ACE engineers visit only some sites. An example of information gleaned from a site visit comes from a sand and gravel plant expansion in Northeast Ohio. An ACE engineer conducting an AJD field visit discovered "small drainage swales" potentially under tree cover connecting to jurisdictional waters and concluded the site was jurisdictional. This example has project identifier LRB-2015-01193.

145 An AJD reports the water resource's coordinates, whether the resource is jurisdictional, the rule under which the AJD was decided, the Cowardin code, the resource type, and a unique project identification code (e.g., different wetlands on a single housing development may have separate AJDs but the same project identifier), which we use to separate AJDs into test and training sets (SM section A.2). AJDs are valid for five years. About half of AJDs also incorporate site visits.

150 ACE requests AJDs to list centroids of water resources. We scrutinized high-resolution satellite imagery around 20 randomly selected AJDs on lakes and found that coordinates for about half of these water resources listed the centroids, while coordinates for the other half listed other locations around the water resource. Due to this focus and the model architecture used, we interpret WOTUS-ML as classifying individual points as regulated or not. Although WOTUS-ML can output different regulatory probabilities inside versus outside water resources, it is better suited at identifying jurisdiction for a water resource overall than delineating wetland boundaries (e.g., ACE sometimes conducts wetland delineation, which seeks to identify the exact boundaries of a wetland feature).

160 During the AJD process, ACE engineers also assign the water resource a Cowardin class and resource type. Cowardin classes are a hierarchical system for classifying wetland and water resources that capture the aquatic system, type of substrate, and water regime of the habitat (14). Resource types correspond to legal groupings of aquatic habitats, such as adjacent wetlands (those abutting a stream/river) or isolated wetlands (those that do not have a surface water connection) specified in each rule. Broad categories of Cowardin classes and resource types relate to different interpretations of WOTUS. For example, *Rapanos* regulates isolated wetlands and NWPR does not.

165 The main text refers to areas where a land use model predicts development. We identify these areas using ICLUS, which provides a raster identifying areas with increasing urban development in the period 2020-2030 (24), which may be more likely to have future AJDs. Using ICLUS, we

170 categorize each point into one of three levels of development. We define a point as undeveloped
if it has any of the following ICLUS categories: natural water; reservoirs; canals; wetlands;
recreation; conservation; timber; grazing; pasture; cropland; mining; or barren land. We define a
point as semi-developed if it has any of the following ICLUS categories: exurban, low; suburban;
175 urban, low; parks; or golf courses. We define a point as developed if it has any of the following
ICLUS categories: exurban, high; urban high; commercial; industrial; institutional; or
transportation. Finally, we define areas with increasing urban development as those that ICLUS
predicts move from undeveloped to semi-developed, semi-developed to developed, or
undeveloped to developed, using the preceding definitions.

180 Several parts of the paper and several datasets discuss isolated wetlands and ephemeral streams.
These water features have different names and sub-types in different regions of the country,
including prairie potholes, vernal pools, Carolina bays, playa lakes, alpine ponds, desert
depressions, swale ponds, and others.

185 WOTUS-ML observes differences across ACE districts and state boundaries. NAIP has some
differences in collection methods across states. For example, the flying season for imagery
collection can differ modestly across states, and photographic methods can also differ (40). The
WOTUS-ML inputs include a state boundary layer, which helps account for differences in NAIP
methodology across states. Differences in regulatory probabilities across ACE districts are salient
in the unprocessed AJD data.

190 The main text reports estimates of the difference in flood mitigation benefits and land values
between *Rapanos* and NWPR. The range we report reflects homogeneous versus heterogeneous
estimates of flood mitigation benefits and land values from other sources. Our homogeneous
estimate uses the finding that the mean US wetland provides \$1,840 to \$1,900 per hectare in
annual flood mitigation benefits (3). Aggregated, this represents \$25 to \$26 billion in annual US
195 flood mitigation benefits, or \$500 to \$520 billion in present value flood mitigation benefits,
discounted at 5%. Heterogeneous estimates of flood mitigation values by population density (3),
applied to our 4 million prediction points, imply that the mean wetland deregulated between
Rapanos and NWPR has annual flood mitigation benefits of \$1030/hectare. Aggregated, this
heterogeneous estimate implies that these deregulated points represent \$12 billion in annual
200 flood mitigation benefits, or \$250 billion in present value. The mean value of US wetlands is
\$12,700/acre (41). Using heterogeneous estimates of land values for each deregulated point (41),
the mean wetland deregulated between *Rapanos* and NWPR has fair market value of
\$8,313/acre. Thus, the heterogeneous estimates imply the national land value of wetlands
deregulated between *Rapanos* and NWPR is \$249 billion. The heterogeneous estimates of flood
205 mitigation benefits and land values for deregulated wetlands are smaller than the homogeneous
estimates for the mean US wetland because deregulated points are in less populated areas than
the mean US point.

210 A.5 Prediction points

For prediction, we collect input data on three distinct samples of points in the CONUS. First, we
collect 4.1 million points at random locations across the entire CONUS. To do this, we first
divide the country into approximately 80,000 0.1 degree by 0.1 degree grid cells. We then
randomly sample 50 points in each cell. We choose sample sizes to support detailed maps and
215 case studies. In practice, input data are not available for about 4% of the points we sample, so we

220 generate predictions for a total of 3.94 million points. The large number of prediction points allows us to produce high-resolution maps, as in Fig. 3, as well as detailed case studies, as in Fig. 4. We additionally use the intersections of these points with areas of particular interest to produce the values in Table 1. In reporting aggregate statistics, the share of the four million prediction points in streams and wetlands that are jurisdictional approximates the share of wetland acres and stream miles that are jurisdictional.

225 The second CONUS sample we collect covers about 3,000 points in a dense sample of approximately 2.7 km sq centered on the Sackett property near the shore of Priest Lake, Idaho. These points enable us to analyze in detail an area of particular interest as it lies as the center of the 2023 Supreme Court *Sackett* case. In addition, these points clarify how spatially precise our predictions are.

230 Finally, we collect approximately 6,200 points along traditional navigable waters legally identified in Title 33 of the US code (23) and link them to corresponding flowline names in NHD. In line with EPA and ACE practice, we use “traditional navigable waters” to denote major waterways primarily used for navigation, sometimes called, “navigable-in-fact waters.” We select these points because they are regulated under every WOTUS rule, and thus are less likely to appear in our AJD training set. These points allow us to test our model’s out of sample
235 performance on locations virtually certain to be jurisdictional.

B Supplementary Text

B.1 Model Accuracy and Potential Uses

240 To evaluate model performance, one metric is whether WOTUS-ML predictions are useful enough for different possible applications. The main text provides a few statistics that help assess the model’s potential usefulness, which we summarize here. For all AJDs, WOTUS-ML test set accuracy is 79%, reflecting learning of 14 percentage points. The corresponding Area Under the Receiver Operating Characteristic Curve (AUC) is 0.85. The ROC curve is plotted in Figure S7.
245 AUC varies little by rule at 0.82, 0.86, and 0.87 for *Rapanos*, CWR, and NWPR respectively. Finally, WOTUS-ML scores provide an unbiased estimate of the probability that a site is regulated. Correspondingly, grid search for the optimal classification threshold returned 0.51. Relative to a threshold of 0.50, using a threshold of 0.51 would increase accuracy on the validation set by 0.013 percentage points (a hundredth of a percentage point); thus, we use a
250 threshold of 0.50.

255 For a developer, learning that a site’s WOTUS-ML score is below 0.07 or above 0.95 provides a strong signal, with over 95% accuracy, whether a site is jurisdictional. Many developers may find this information useful for investment and permitting decisions. For example, observing the WOTUS-ML prediction that a site is 95% likely to be regulated may lead a developer to request a PJD rather than an AJD, which can increase permitting efficiency. Sites with WOTUS-ML scores closer to 0.5 provide weaker signals for an individual site, since accuracy is lower for those score ranges, but may still provide useful inputs for some developer investment and permitting decisions. For ACE, having WOTUS-ML scores available could help to prioritize between sites where WOTUS-ML indicates jurisdiction is more certain, so the return to
260 additional ACE effort is lower (or a field visit may be less needed), versus sites where WOTUS-ML scores are closer to 0.5, and where the return to ACE effort may be greater. WOTUS-ML

could flag cases where an ACE decision differs substantially from the WOTUS-ML prediction, and which ACE could use to select cases for review. Additionally, WOTUS-ML scores could allow benchmarking or provide a training tool for new ACE engineers or provide a way of comparing practices across ACE districts. Under current practice, algorithms like WOTUS-ML alone cannot provide a legally-binding jurisdictional determination, but could provide a useful input to ACE's decision process.

For EPA, environmental, or industry associations, WOTUS-ML could help estimate the share of waters that are regulated under different rules, geographic areas, or water types. The unbiasedness described in Figure 2A is useful here, since it shows that while WOTUS-ML has imperfect accuracy for individual sites, these errors average out in large samples.

Another metric to evaluate model performance is whether WOTUS-ML exceeds performance of available alternative tools. We are unaware, however, of any available alternative tools. We believe no other analyst in government, academia, or the private sector has constructed a model that predicts AJDs and has then evaluated the accuracy of this model. In this sense, a possible use of WOTUS-ML is to highlight the importance of this prediction problem and to provide a set of accuracy statistics upon which other, potential models may attempt to improve.

We briefly discuss how WOTUS-ML differs between sites with and without a field visit. Under COVID-19 lockdowns, some sites had a "visit, conducted virtually using remote tools," primarily under NWPR, which we count as not having a site visit. The main text notes that test set accuracy is 82% for AJDs without a field visit and 74% for AJDs with a field visit. The share regulated is 26% for AJDs without a field visit and 46% for AJDs with a field visit. Thus, learning is only 8 percentage points for sites without a field visit, but 20 percentage points for sites with a field visit. At the same time, the low regulatory probability for AJDs without a field visit overall (26%) gives less scope for model learning.

Regulation and model learning differ for AJDs with a field visit for several reasons. On a field visit, ACE engineers can discover evidence of regulated waters that computer analysis misses, such as drainage swales under forest cover, plant species indicative of wetlands, or the location of the ordinary high-water mark. Additionally, much of the time when NWPR was operating occurred in 2020 and 2021 when travel was limited due to COVID-19. Hence field visits occurred more under *Rapanos* and CWR than under NWPR, and *Rapanos* and CWR regulate a larger share of waters. Finally, ACE engineers choose field visits purposefully, for more complex cases (39).

WOTUS-ML has imperfect accuracy in part due to characteristics of the training data. AJDs (the training data) can have human error or reflect brief and uncommon environmental conditions (42, 43). Implementation varies; for example, ground-truth data reveal sharp regulatory differences around ACE district boundaries (Fig. S2), and individual engineers and field offices could vary further. After an AJD is finalized, landowners receive a form to request an appeal (44); the existence of this form implicitly acknowledges that AJDs can be incorrect. In-person visits for roughly half of AJDs provide information not available to train WOTUS-ML. In addition, most AJD PDFs note that the decision reflects maps, data, photos, or other information provided by the applicant or an associated environmental consultant about the site, information which is unavailable to train WOTUS-ML.

305 The main text explains how the permutation tests highlight which features most influence the
model’s predicted probabilities. It is also informative to observe which features play less of a
role. ACE documentation notes that soils, hydrology, and vegetation determination can play an
important role in identifying wetlands (21). Our gNATSGO extract includes five types of
310 categorical variables: soil taxonomic class; soil hydric rating; flooding frequency; ponding
frequency; and water table depth variables, in addition to others (Table S3). Fig. S10 shows that
gNATSGO is moderately important nationally and is the third-most important feature to
determine which sites are jurisdictional within a state. For reasons discussed in SM section A.1,
we permute features in groups, so we identify the importance of gNATSGO as a whole, but not
the individual features of gNATSGO. Nonetheless, this would suggest that soil and hydrology
315 variables play an important role in guiding WOTUS-ML’s predictions across sites within a state.
One can similarly look at other permutation importance levels for Fig. S10, compare them to the
individual variables listed in Table S3, and determine their relative importance levels.

320 B.2 Data Sources

To compare our data against the data sources that ACE engineers consult, we selected a random
sample of 20 AJDs, obtained the PDF decisions for these AJDs from ACE’s ORM2 online
database, and categorized the data sources used in these decisions.² The five most commonly-
used data sources were maps and data sheets submitted by the applicant and associated
325 consultant (100% of sampled AJDs), aerial imagery (100%), topographic maps (90%), NWI
(75%), and soil surveys (65%). The AJDs also report that other data sources are less widely used,
including ACE’s Antecedent Precipitation Tool (30%). HUC maps (25%), NHD (20%), LiDAR
(20%), state/local flood maps (15%), FERA/FIRM maps (10%), ACE data sheets (5%), and ACE
navigable waters studies (5%). Apart from the applicant materials, which have unspecified data
330 sources, our data cover aerial imagery via NAIP, topographic maps via the DEM, NWI, and soil
surveys via gNATSGO.

Observing that many AJDs use a dataset does not indicate that the dataset most influences
decisions. Nonetheless, we note that NWI and gNATSGO are the most important for WOTUS-
335 ML decisions within a state in Fig. S10, and are among the most commonly used data sources in
these AJDs. We also note the contrast that aerial imagery is used in 100% of these AJDs, but
NAIP is much less important in Fig. S10. One potential reason for this is that much of the
information in NAIP may already be in our other datasets, some of which ACE does not directly
consult (e.g., ecoregions, PRISM, and NLCD).

340 B.3 Decision Model and Cost Savings

WOTUS-ML can support several parts of the CWA Section 404 process. At decision (a) in Fig.
S1, WOTUS-ML scores can give the developer information that helps decide whether to hire a
345 consultant, develop while ignoring WOTUS, or stop development. At (b), WOTUS-ML could
give ACE and EPA potential signals for prioritizing potential enforcement cases. At (c),
WOTUS-ML could give a developer and consultant useful information about deciding between
PJDs, AJDs, development, or non-development. At (d), WOTUS-ML as a decision support tool
could give ACE useful information about the estimated probability an AJD is regulated.

² This sample excluded AJDs missing PDFs or AJDs corresponding to more than one PDF in ORM2.

350 We can use this cost to provide an estimate of one component of WOTUS-ML's value, at
decision node (c). Because this only includes one component of WOTUS-ML's informational
value at this decision node, and since WOTUS-ML has potential value at other decision nodes, it
may provide a lower bound on the value of WOTUS-ML to stakeholders.

355 Specifically, Fig. 2A shows that for 27% of AJDs, WOTUS-ML provides greater than 95%
certainty about whether a site is jurisdictional. For these sites, requesting an AJD is more likely
to be an error, since a PJD or development without permit are alternatives. The expected cost of
the error depends on the number of decisions at each part of the tree and the costs of the other
components, most of which are unknown. In the absence of this data, we can obtain one ballpark
360 number from EPA and ACE's (2) estimate that each Section 404 permit costs \$5,000 to \$39,000,
representing S in the decision tree. We are unaware of estimates for the cost of AJDs (A), but
these estimates of the permit (S) cost provide a plausible starting estimate. If anything, AJD costs
might exceed this amount since an environmental consultant alone can exceed this amount, and
then AJDs create additional costs due to delay, uncertainty, and resulting changes to project
design.

365 B.4 External Validity

This paper asks what the CWA regulates, using data over the period 2015-2022. In this sense,
predicting CWA jurisdiction in distant future years is not our primary focus. At the same time,
for stakeholders, it may be useful to understand the external validity of WOTUS-ML to *Sackett*
or future regulation. Given that WOTUS-ML inputs include non-geophysical layers like state
370 boundaries, ACE district boundaries, and distance to a field office, it is unclear how district-
specific policies could change over time. The state boundaries with the sharpest regulatory
discontinuities in maps like Fig. 3A also lie on ACE district boundaries. Except for the few states
which have deregulated AJD determination, our sense is that ACE districts rather than states
provide the most important heterogeneity. Additionally, the importance of PRISM climate data
375 and our use of 30-year climate normals could mean that climate change would affect future
predictive accuracy.

We believe that five and possibly ten future years is the most relevant period for stakeholders. In
part this is because AJDs are valid for five years, because a developer who requests an AJD is
likely to develop promptly, and because CWA rules in the last decade have changed frequently.
380 For policymakers, federal budget scoring often considers ten years.

Our cautious assessment is that conditional on rule changes, WOTUS-ML is likely to have good
external validity for five to ten years. Given the new *Sackett* rule beginning in late 2023, one
could train WOTUS-ML in 2024 or later on *Sackett* AJDs, and then we would expect similar
385 external validity. External validity is more difficult to assess beyond that period. The WOTUS-
ML training data covers the eight-year period from 2015 to 2022, which is a similar duration as a
five- to ten- year prediction. The role of PRISM in prediction occurs despite the presence of
climate change in this period, and our sense is that the PRISM data are highlighting cross-
sectional variation in deserts, moisture, heat, and other features that will be broadly similar in the
390 next 5-10 years. If ACE district jurisdictional practices or climate substantially changed after
several future years, one could always train a new version of WOTUS-ML on newer data or
update the model inputs, although that possibility does not shed light on the relevance of the
current WOTUS-ML for predicting future jurisdiction.

395 We have asked policymakers why ACE district practices differ, since answering this question
could help assess the persistence of these patterns. We do not have a single clear answer. While
ACE seeks to harmonize AJD criteria across districts, ACE regulatory practices are more
400 decentralized than some federal agencies. Some ACE districts and regions have supplements to
the ACE handbook that guide specific aspects of AJDs for local terrain. We also emphasize that
ACE district staff are not federal political appointees, and while ACE district leaders have
influence, one might expect less variation in ACE district priorities over time than occurs among
some federal agencies where presidential appointees change regularly.

405 We also train a separate deep learning model, AJD-ML, which helps us study WOTUS-ML's
external validity. AJD-ML predicts the probability that an arbitrary location has an AJD. By
contrast, WOTUS-ML predicts the probability that an AJD for a location concludes that the
location has jurisdictional waters. In other words, AJD-ML and WOTUS-ML have two main
differences: AJD-ML predicts the probability a point has an AJD while WOTUS-ML predicts
410 the probability that an AJD is jurisdictional; and AJD-ML is trained on both AJDs and non-AJD
locations, while WOTUS-ML is only trained on AJDs.

Apart from these differences, AJD-ML and WOTUS-ML are similar. They have identical model
architecture, training, and train-validation-test split procedure. The input data are identical except
415 that AJD-ML does not include a WOTUS rule layer. In other words, to predict which points have
AJDs in AJD-ML, we use aerial imagery, maps of streams and wetlands, soil quality and
groundwater information, a digital elevation model, land use data, etc. To train and evaluate
AJD-ML, we use all AJD points and 152,201 of our four million prediction points. We randomly
choose these prediction points, subject to not overlapping with any AJD. We train AJD-ML for
420 11 epochs. AJD-ML's test set accuracy is 92.7%.

Fig. S8 compares WOTUS-ML accuracy across deciles of AJD-ML probabilities. In other
words, it investigates how accurately we can predict whether a given location is jurisdictional
under the CWA (WOTUS-ML model), separately for points that are more versus less likely to
425 have an AJD (AJD-ML model). In the top panel, the horizontal axis shows bins indexing the
probability that a given point has an AJD. These bins represent fitted probabilities from the AJD-
ML model. The vertical axis shows WOTUS-ML's accuracy in each bin (blue triangles),
measured using all AJDs; and the vertical axis also shows the share of AJDs in each bin that
ACE determines to be jurisdictional (green circles).

430 Our main finding from Fig. S8 is that WOTUS-ML has similar accuracy of about 80% for both
locations that look very much and very little like a typical AJD. We obtain this finding from the
observation that the blue triangles in the top panel have flat slope horizontally, at a value near 0.8
on the y-axis. WOTUS-ML's accuracy is similar for sites that have characteristics typical of
AJDs, on the graph's right side, as for sites that have characteristics atypical for AJDs, on the
435 graph's left side.

This finding that WOTUS-ML's accuracy is similar for points that do and do not look like a
typical AJD helps increase confidence in our headline conclusions about the share of all US
440 wetlands and streams that are jurisdictional. One might have the concern that the wetlands and
stream sites which have AJDs are larger, more complex, more ambiguous, or for some other
reason different than the mean US wetland or stream. This concern could imply that WOTUS-

ML has good accuracy on the kinds of sites on which WOTUS-ML is mostly trained – sites that look like a typical AJD – but not on other kinds of sites. This concern could therefore imply that our statistics on accuracy and jurisdiction of WOTUS-ML might not extrapolate well to all US streams and wetlands. Fig. S8 does not support this concern, and instead suggests that WOTUS-ML has similar accuracy for all types of points, and not only those that look like a typical AJD. It therefore suggests that we can reasonably use WOTUS-ML to assess the share of all streams and wetlands nationally that are jurisdictional, even including streams and wetlands with different characteristics than most AJDs.

B.5 Sackett

In May 2023, the Supreme Court issued a decision in *Sackett* (45). In August 2023, the EPA issued a final rule which conforms to *Sackett* (46). *Sackett* strikes down the significant nexus standard, and excludes interstate wetlands and wetlands separated from jurisdictional waters by a man-made dike or barrier, natural river berm, beach dune, or the like. Formally, the EPA’s August 2023 rule modifies the January 2023 rule issued under the Biden administration, which is similar to *Rapanos*, rather than modifying NWPR. *Sackett* and the *Sackett* rule focus on changing the significant nexus standard but focus less on the relatively permanent standard.

We here discuss two potential ways to quantify what *Sackett* regulates. This paper trains a machine learning model on past AJDs issued under a given rule. This is infeasible for *Sackett* in the near term since the EPA has just issued the *Sackett* rule. It will take time for many *Sackett* AJDs to be issued. The recent history of CWA regulation also has had numerous delays and obstacles to changing CWA rules and implementing new rules. It is possible that procedural delays could slow implementation of the *Sackett* rule.

A possible alternative strategy to quantify what *Sackett* regulates, which differs from the methodology applied in the rest of the paper, would be a relabeling technique. Because the EPA’s *Sackett* rule modifies the January 2023 rule, which is very similar to *Rapanos*, this methodology would identify which past AJDs already completed under *Rapanos* would lose jurisdiction under *Sackett*. This relabeling technique would then construct a synthetic dataset of *Sackett* AJDs by changing the labels (the jurisdictional outcomes) for the AJDs that would lose jurisdiction in *Sackett* relative to *Rapanos*. Applying this relabeling methodology would then train WOTUS-ML on this relabeled set of AJDs, and predict jurisdiction for the 4 million and other points. These points would describe a prediction of what *Sackett* regulates.

While we investigated the potential of this relabeling methodology in detail, we do not implement it here because we believe that without *Sackett* AJDs for training a model, it is difficult to implement it with sufficient accuracy. Specifically, the construction of the *Sackett* rule makes it difficult to conclusively identify which *Rapanos* AJDs would lose jurisdiction under *Sackett*. The *Sackett* rule removes the significant nexus standard for several legal categories of waters identified in the January 2023 rule. The legal categories of waters identified in *Rapanos* AJDs (formally, called “resource types” in the AJD data) differ from the legal categories of waters listed in the January 2023 rule. Thus, determining which *Rapanos* AJDs would lose jurisdiction under *Sackett* depends on a match in legal water categories between the two rules. More importantly, for several legal categories of waters in the January 2023 rule, *Sackett*’s removal of the significant nexus standard deregulates some but not all waters in the category. Within a category, the *Rapanos* AJDs alone have insufficient information both about what share and which AJDs would lose jurisdiction under *Rapanos*.






Fig. S1: A Decision Tree Clarifies the Process of Development and Jurisdictional Determinations and Shows Potential Uses of WOTUS-ML. Each box describes an intermediate decision and each oval describes a potential terminal outcome. Branches and decision nodes are in blue and lowercase, costs and payoffs are in capital red. At (a), the developer chooses whether to develop while ignoring the CWA, hire an environmental consultant to support the CWA Section 404 process, or not develop. If the project proceeds assuming jurisdictional Waters of the United States (WOTUS) are not present, at (b), ACE and EPA may discover the development, conclude WOTUS are present, and require restoring the site to its original conditions and imposing a fine. At (c), the developer and an environmental consultant may provide a PJD, request an AJD, proceed ignoring WOTUS, or not develop. At (d), ACE determines whether the AJD is WOTUS. If the AJD is not WOTUS, development occurs. At (e), if the AJD is WOTUS, the developer may obtain a Section 404 permit and develop or may stop development. AJD are approved jurisdictional determinations, PJD are preliminary jurisdictional determinations, and WOTUS are Waters of the United States. The model shows common sequences but abstracts from others (e.g., a developer could provide a PJD without a consultant, and might provide a PJD but not develop the project). Payoff variables are follows: C is cost of hiring an environmental consultant; P is cost of obtaining a PJD, including paperwork, delays, and uncertainty; A is cost of obtaining an AJD, including paperwork, delays, and uncertainty; S is cost of obtaining and complying with a Section 404 nationwide or individual permit, including mitigation, compensatory measures, or constraints on development; D is payoff from development; and V is fines and other costs due to violating CWA Section 404.


A AJs, all rules



B AJs, *Rapanos*



C AJs, CWR



D AJDs, NWPR




Fig. S2: AJDs (Training Data), by Rule. (A) All AJDs. (B) AJDs decided under *Rapanos*. (C) AJDs decided under CWR. (D) AJDs decided under NWPR. Green dots represent jurisdictional sites; brown dots represent non-jurisdictional sites.





Fig. S3: Train, validation and test assignment. Train points are in green, validation points in brown, and test points in blue. Black lines show ACE district boundaries. We plot the validation and test sets on top of the train set.


A National Agricultural Imagery Program




B National Hydrography Dataset




C Gridded National Soil Survey
Geographic Database



D National Wetlands Inventory



E National Land Cover Database



F PRISM




Fig. S4: Maps of six input layers. (A) shows NAIP imagery. (B) shows NHD flowlines, where darker colors represent streams that flow more directly into oceans or great lakes (stream order). (C) shows soil taxonomic class from gNATSGO. (D) shows NWI polygons. (E) shows land cover. (F) shows precipitation patterns from PRISM.






Fig. S5. Sample Inputs for One Site. Fig S5 depicts the 34 input layers for WOTUS-ML (see Table S3 for details). The location shown here is associated with an Approved Jurisdictional Determination (AJD) located at -121.795E, 39.734N in Butte County, California. The nearest water body is Dead Horse Slough, which flows into the Traditional Navigable Water (TNW) Little Chico Creek. The National Agriculture Imagery Program (NAIP) imagery shows that the site

includes an undeveloped lot besides a waterbody with adjacent wetlands. The National Wetlands Inventory (NWI) shows three types of wetlands at this location: Riverine (light blue), Freshwater Emergent Wetlands (light green), and Freshwater Forested/Shrub Wetland (dark green). The National Land Cover Database (NLCD) layer includes open space (pink), developed land (red), and grasslands (tan). 3DEP shows that elevation ranges from 73m in the streambed to 78m in the field. The water feature appears in the National Hydrography Dataset (NHD) layer as an intermittent stream (FCode 46003). WOTUS-ML also inputs soil characteristics in the gridded National Soil Survey Gridded Database (gNATSGO) layers, information on climate in the PRISM layers, and geographic variables such as state, ecoregion, and U.S. Army Corps of Engineers (ACE) district. Finally, the model includes a one-hot encoding for each WOTUS rule: *Rapanos*, the Clean Water Rule (CWR), or the Navigable Waters Protection Rule (NWPR). This site was determined to be jurisdictional under *Rapanos*. WOTUS-ML assigns this site a score (i.e., probability of being jurisdictional) of 96% under *Rapanos*.


A *Rapanos*, AJD Test Set




B *Rapanos*, Four Million Prediction Points




C CWR, AJD Test Set



D CWR, Four Million Prediction Points



E NWPR, AJD Test Set



F NWPR, Four Million Prediction Points




Fig. S6: Histograms of WOTUS-ML Scores for AJD Test Set and Four Million Prediction Points, By Rule.




Fig S7: WOTUS-ML achieves an AUC of 0.85. The Receiver Operator Characteristic (ROC) curve is plotted by calculating false positive and false negative rates for various classification thresholds. The Area Under the ROC Curve (AUC) is the integral of the ROC curve along the x-axis.






Fig. S8: WOTUS-ML Accuracy is Not Correlated with Predicted AJD Probability. Predicted AJD probabilities are merged with WOTUS-ML predictions, then WOTUS-ML accuracy is calculated for each 10 percentage point bin of predicted AJD probability. The WOTUS-ML accuracy for each bin is plotted as a blue triangle. We find no correlation between WOTUS-ML accuracy and predicted AJD probability. We also plot the share of AJDs in each bin that are WOTUS as green circles. We find a weak positive correlation between the predicted probability of being an AJD and the probability of being WOTUS. The bottom panel shows histograms of the distribution of predicted AJD probabilities for the AJDs and the sampled prediction points. Note that this figure includes points from all folds (training, validation, and test) to maximize the number of observations in each bin. The test set only (not shown) has a similar horizontal pattern, though with more variance given the smaller sample. This has little effect on interpretation because WOTUS-ML and AJD-ML both exhibit minimal overfitting.


A Left Tail Accuracy



B Right Tail Accuracy



C Left Tail Share



D Right Tail Share




Fig. S9: Cumulative Shares and Accuracy in AJD Test Set. The x-axis in (A) lists threshold predicted regulatory probabilities such that for test set AJDs below each threshold, WOTUS-ML has accuracy of 99%, 95%, 90%, or 85%. (B) is similar but calculates accuracy for AJDs above each threshold. (C) calculates the cumulative share of test set AJDs in the test set with predicted regulatory probability below the thresholds identified in (A). (D) is similar but calculates the cumulative share of test set AJDs above each threshold. Summing the y-axis values in (C) and (D) gives the y-axis in Fig. 2A.




Fig. S10: Permutation Tests of WOTUS-ML Feature Importance. Features are permuted across samples in validation set (A) nationally and (B) within each state. The datasets represent the following information: NAIP is imagery, NWI and NHD are wetlands and streams, DEM is elevation, NLCD is land cover, PRISM is climate, and gNATSGO is soil and groundwater.

Table S1: Categorization of Nine Hydrological (Cowardin) Codes

Cowardin code	Description
1. Streams, ephemeral	A wetland, spring, stream, river, pond or lake that only exists for a short period
2. Streams, intermittent	Intermittent, Riverine; Streambed, Intermittent, Riverine
3. Streams, perennial and other	Upper Perennial, Riverine; Lower Perennial, Riverine; Unknown Perennial, Riverine
4. Wetland, emergent	Emergent, Palustrine; Persistent, Emergent, Palustrine; Nonpersistent, Emergent, Palustrine
5. Wetland, forested	Forested, Palustrine; Broad-Leaved Deciduous, Forested, Palustrine; Needle-Leaved Evergreen, Forested, Palustrine"; Needle-Leaved Deciduous, Forested, Palustrine; Broad-Leaved Evergreen, Forested, Palustrine; Indeterminate Deciduous, Forested, Palustrine
6. Wetland, other	All other palustrine
7. Estuaries	Estuarine
8. Uplands	Uplands
9. Other	Marine, lacustrine, riparian

Table S2: Categorization of Nine Resource Types. RPW, relatively permanent waters; TNW, territorial national waters.

	<i>Rapanos</i>	CWR	NWPR
1. Ephemeral			(b)(3) Ephemeral feature, including an ephemeral stream, swale, gully, rill, or pool
2. Isolated	Isolated (interstate or intrastate) waters		(b)(1)
3. Non-RPW that flows directly or indirectly into TNW	Non-RPW that flows directly or indirectly into TNW		
4. Other non-jurisdictional		(b)(1), (b)(2), (b)(3)(i), (b)(3)(ii), (b)(3)(iii), (b)(4)(v), (b)(5), (b)(7), (b)(4)(iii), (b)(4)(iv), (b)(4)(ii), (b)(4)(i), (b)(4)(vi), (b)(6), (b)(4)(vii)	(b)(2), (b)(4), (b)(5), (b)(6), (b)(7), (b)(8), (b)(9), (b)(10), (b)(11), (b)(12), the review area is comprised entirely of dry land
5. Other jurisdictional		(a)(4), (a)(7), (a)(8)	(a)(3)
6. RPW that flows directly or indirectly into TNW	Relatively Permanent Water that flows directly or indirectly into Traditional Navigable Water	(a)(5)	(a)(2)
7. TNW	Traditional Navigable Water	(a)(1), (a)(2)	(a)(1)
8. Uplands	Uplands	Uplands	
9. Wetlands and adjacent/abutting regulated waters	Wetlands Directly Abutting RPW that flows directly or indirectly into TNW; Wetland Adjacent to Non-RPW that flows directly or indirectly into TNW; Wetlands Adjacent but not Directly Abutting RPW that flows directly or indirectly into TNW; Wetlands Adjacent to TNW	(a)(6)	(a)(4)

Table S3: Datasets to Support Predictions of Jurisdictional Waters.

Dataset	Input layer	Variable definition	Data type	Spatial Resolution	Source
National Agriculture Imagery Program (NAIP)	Red band	Red channel visible light	Raster	0.6 to 1.0 meters	(15)
	Blue	Blue channel visible light			
	Green	Green channel visible light			
	NIR	Near-infrared light			
National Wetlands Inventory (NWI)	Wetland type	NWI wetland types: Estuarine and Marine Deepwater, Estuarine and Marine Wetland, Freshwater Emergent Wetland, Freshwater Forested/Shrub Wetland, Freshwater Pond, Lake, Riverine, Other	Vector	1:250,000	(19)
National Hydrography Dataset (NHD) Plus V2	FCode	Water feature type (e.g. perennial stream, intermittent stream, coastline)	Vector	1:100,000	(17)
	Path length	NHD flowline distance			
	High flow	Maximum flow for this water segment over a sequential 3-month period, using NHD Value Added Attributes Enhanced Runoff Method (EROM) long-term mean flow estimates for each month.			
	Low flow	Minimum flow for this water segment over a sequential 3-month period, using EROM.			
	Stream order	Hierarchy of streams from the source (or headwaters) downstream			
USGS 3-Dimensional Elevation Program (3DEP)	Elevation	Height above sea-level	Raster	10 meters	(30)
U.S. EPA Ecoregions	Level IV Ecoregion	Ecoregions are areas where ecosystems (and the type, quality, and quantity of natural resources) are generally similar. There are 967 level IV ecoregions in the United States.	Vector	1:250,000	(31)

Continued next page

Dataset	Input layer	Variable definition	Data type	Spatial Resolution	Source
Parameter-elevation Regressions on Independent Slopes Model (PRISM) 30-year Normals	Precipitation	Average annual total precipitation	Raster	4 kilometers	(32)
	Minimum temperature	Daily minimum temperature, averaged over 1990-2021			
	Maximum temperature	Daily maximum temperature, averaged over 1990-2021			
	Mean temperature	Daily mean temperature, averaged over 1990-2021			
	Mean dew point temperature	Daily mean dew point temperature (the temperature to which air must be cooled to become saturated with water vapor), averaged over 1990-2021			
	Minimum vapor pressure deficit (VPD)	Minimum VPD (difference between the amount of moisture in the air and how much moisture the air can hold), averaged over 1990-2021			
	Maximum vapor pressure deficit (VPD)	Maximum VPD (difference between the amount of moisture in the air and how much moisture the air can hold), averaged over 1990-2021			
	Solar radiation (clear sky)	Total daily global shortwave solar radiation received on a horizontal surface, averaged over 1990-2021			
	Solar radiation (total)	Total solar radiation incident on a horizontal surface), averaged over 1990-2021			
	Cloudiness	Atmospheric transmittance (cloudiness), averaged over 1990-2021			
U.S. Army Corps Regulatory Boundaries	District codes	Each ACE district is assigned a unique value.	Point	1:250,000	(20)
	Distance to headquarters	We calculate the distance from each point to the district headquarters.			

Continued next page

Dataset	Input layer	Variable definition	Data type	Spatial Resolution	Source
Gridded National Soil Survey Geographic Database (gNATSGO)	Taxonomic class	The Soil Taxonomy subgroup and family for a soil.	Raster	30 meters	(16)
	Hydric rating	Is the map unit “hydric soil”?			
	Flooding frequency	The annual probability of a flood event expressed as a class.			
	Ponding frequency	The number of times ponding occurs over a year			
	Water table depth	The shallowest depth to a wet soil layer			
National Land Cover Database (NLCD)	Landcover	The NLCD has 20 land cover classes: Open water, ice/snow, four classes of developed land (open, low, medium, and high), barren, three forest classes (evergreen, deciduous, mixed), two scrub classes (dwarf, shrub), four herbaceous classes (grassland, sedge, moss, lichen), two agricultural classes (pasture/hay, cultivated), and two wetland classes (woody, emergent herbaceous)	Raster	30 meters	(33)
Topologically Integrated Geographic Encoding and Referencing System (TIGER)/Line State boundaries	State codes	Each state is assigned a unique value	Vector	1:250,000	(34)
CWA Approved Jurisdictional Determinations	WOTUS rule	Three WOTUS rules: <i>Rapanos</i> , CWR, NWPR	Point	--	(38)

Table S4: Accuracy of WOTUS-ML Model Predictions, All Rules. Statistics describe the AJD test set. Column (1) shows the share that USACE determines to be regulated. Column (2) shows the share that WOTUS-ML predicts to have score above 0.5. Column (3) shows the share that WOTUS-ML correctly classifies as regulated. Column (4) shows the number of test set AJDs. Field visit indicators and Cowardin codes are from the AJD data. NWI and NHD include points within 5-10 meters of these vector data. **(E)** shows five ACE districts selected to represent districts with among the largest or smallest numbers of AJDs, and to show geographically heterogeneous districts.

	AJDs: regulated (1)	WOTUS-ML: score > 0.5 (2)	Accuracy (3)	N (4)
A General groups of points				
All AJDs	0.35	0.29	0.79	15,970
USACE field visit	0.46	0.38	0.74	7,198
USACE no field visit	0.26	0.21	0.82	8,772
B By Cowardin Codes				
All rivers and streams (riverine)	0.43	0.31	0.78	4,353
Wetlands (palustrine)	0.38	0.28	0.77	8,203
Uplands	0.00	0.17	0.83	2,529
Estuaries (estuarine)	0.99	0.94	0.94	304
Lakes (lacustrine)	0.39	0.30	0.81	352
C By Vector Data Overlay				
All rivers and streams (NHD)	0.71	0.75	0.77	768
All wetlands (NWI)	0.53	0.45	0.75	2,247
D By year				
2016	0.52	0.46	0.77	1,731
2017	0.41	0.40	0.82	1,255
2018	0.34	0.32	0.80	1,580
2019	0.38	0.37	0.76	2,067
2020	0.31	0.22	0.79	3,816
2021	0.25	0.16	0.79	3,982
2022	0.43	0.39	0.75	918
E Selected USACE Districts				
Albuquerque	0.13	0.05	0.86	100
Charleston	0.51	0.44	0.73	935
Philadelphia	0.62	0.69	0.69	124
St. Paul	0.02	0.01	0.98	1,296
Wilmington	0.78	0.84	0.79	868

Table S5. Confusion Matrix. For each panel, table entries show share of test set AJDs in each cell. Top-left cell within a panel represents true positives, bottom-right represents true negatives, and top-right and bottom-left represent false positives and false negatives. Share of true positives plus true negatives equals accuracy.

		AJDs: Test set outcomes	
		Regulated (1)	Not regulated (2)
A All rules			
WOTUS-ML Prediction			
	Regulated	0.22	0.08
	Not regulated	0.14	0.57
B Rapanos			
WOTUS-ML Prediction			
	Regulated	0.29	0.09
	Not regulated	0.13	0.50
C CWR			
WOTUS-ML Prediction			
	Regulated	0.29	0.12
	Not regulated	0.11	0.48
D NWPR			
WOTUS-ML Prediction			
	Regulated	0.11	0.05
	Not regulated	0.16	0.69

Table S6: Accuracy of WOTUS-ML Model Predictions, *Rapanos*. Statistics describe the AJD test set. Column (1) shows the share that USACE determines to be regulated. Column (2) shows the share that WOTUS-ML predicts to have score above 0.5. Column (3) shows the share that WOTUS-ML correctly classifies as regulated. Column (4) shows the number of test set AJDs. Field visit indicators and Cowardin codes are from the AJD data. NWI and NHD include points within 5-10 meters of these vector data. (E) shows five ACE districts selected to represent districts with among the largest or smallest numbers of AJDs, and to show geographically heterogeneous districts.

	AJDs: regulated	WOTUS-ML: score > 0.5	Accuracy	N
	(1)	(2)	(3)	(4)
A General groups of points				
All AJDs	0.41	0.37	0.78	8,198
USACE field visit	0.53	0.49	0.76	4,011
USACE no field visit	0.30	0.26	0.80	4,187
B By Cowardin Codes				
All rivers and streams (riverine)	0.67	0.54	0.80	1,425
Wetlands (palustrine)	0.47	0.36	0.76	4,199
Uplands	0.00	0.19	0.81	2,059
Estuaries (estuarine)	1.00	0.96	0.96	236
Lakes (lacustrine)	0.59	0.49	0.77	156
C By Vector Data Overlay				
All rivers and streams (NHD)	0.79	0.87	0.84	370
All wetlands (NWI)	0.61	0.55	0.76	1,285
D By year				
2016	0.52	0.46	0.77	1,731
2017	0.41	0.40	0.82	1,255
2018	0.32	0.30	0.83	1,260
2019	0.39	0.34	0.75	1,217
2020	0.38	0.35	0.82	902
2021	0.35	0.31	0.71	597
2022	0.43	0.39	0.75	918
E Selected USACE Districts				
Albuquerque	0.26	0.11	0.72	47
Charleston	0.65	0.66	0.77	542
Philadelphia	0.48	0.79	0.69	42
St. Paul	0.04	0.01	0.97	618
Wilmington	0.85	0.93	0.84	610

Table S7: Accuracy of WOTUS-ML Model Predictions, NWPR. Statistics describe the AJD test set. Column (1) shows the share that USACE determines to be regulated. Column (2) shows the share that WOTUS-ML predicts to have score above 0.5. Column (3) shows the share that WOTUS-ML correctly classifies as regulated. Column (4) shows the number of test set AJDs. Field visit indicators and Cowardin codes are from the AJD data. NWI and NHD include points within 5-10 meters of these vector data. (E) shows five ACE districts selected to represent districts with among the largest or smallest numbers of AJDs, and to show geographically heterogeneous districts.

	AJDs: regulated	WOTUS-ML: score > 0.5	Accuracy	N
	(1)	(2)	(3)	(4)
A General groups of points				
All AJDs	0.26	0.15	0.79	6299
USACE field visit	0.33	0.16	0.71	2478
USACE no field visit	0.21	0.15	0.85	3821
B By Cowardin Codes				
Rivers and streams (riverine)	0.30	0.15	0.77	2467
Wetlands (palustrine)	0.26	0.16	0.80	3345
Uplands	0.00	0.09	0.91	240
Estuaries (estuarine)	0.88	0.79	0.73	33
Lakes (lacustrine)	0.10	0.05	0.90	129
C By Vector Data Overlay				
All rivers and streams (NHD)	0.60	0.60	0.68	329
All wetlands (NWI)	0.38	0.24	0.73	798
D By year				
2020	0.29	0.18	0.78	2914
2021	0.24	0.13	0.80	3385
E Selected USACE Districts				
Albuquerque	0.00	0.00	1.00	46
Charleston	0.33	0.14	0.67	366
Philadelphia	0.63	0.47	0.67	49
St. Paul	0.02	0.00	0.98	479
Wilmington	0.54	0.56	0.61	208

Table S8: Accuracy of Cowardin Code and Resource Type Predictions. Accuracy takes test AJDs that WOTUS-ML classifies in a particular category (e.g., Cowardin code of ephemeral streams) and reports the share of these AJDs where the ACE engineers classifies the resource as in this category.

	WOTUS-ML:			
	AJDs: share	score > 0.5	Accuracy	N
	(1)	(2)	(3)	(4)
A. Cowardin Codes				
1. Streams, ephemeral	0.14	0.18	0.66	2,265
2. Streams, intermittent	0.07	0.03	0.11	1,128
3. Streams, perennial and other	0.06	0.02	0.16	960
4. Wetland, emergent	0.29	0.34	0.65	4,655
5. Wetland, forested	0.16	0.19	0.61	2,477
6. Wetland, other	0.07	0.01	0.05	1,071
7. Estuaries	0.02	0.03	0.85	304
8. Uplands	0.16	0.18	0.63	2,529
9. Other	0.04	0.02	0.13	581
Overall	1.00	1.00	0.52	15,970
B. Resource Types				
1. Ephemeral	0.09	0.11	0.75	1,388
2. Isolated	0.29	0.36	0.70	4,664
3. Non-RPW that flows directly or indirectly into TNW	0.03	0.01	0.26	424
4. Other non-jurisdictional	0.13	0.11	0.53	2,070
5. Other jurisdictional	0.01	0.01	0.12	164
6. RPW that flows directly or indirectly into TNW	0.11	0.06	0.20	1,698
7. TNW	0.02	0.03	0.67	353
8. Uplands	0.13	0.12	0.57	2,066
9. Wetlands adjacent/abutting regulated waters	0.20	0.18	0.44	3,143
Overall	1.00	1.00	0.54	15,970

Table S9: Regulated Stream Miles and Wetland Acres, by State. Total stream miles in (2) is from NHD stream and river flowline features. Total wetland acres in (3) is from NWI. Regulation rates in (4), (5), (6), and (7) are from WOTUS-ML, applied to the subset of four million prediction points that are within 10 meters of NHD or NWI features. The difference in column (6) is measured in stream miles, and in column (9) in wetland acres. Positive entries in (6) are associated with perennial streams, see SM section A.4.

State (1)	Total Stream miles (2)	Total Wetland Acres (3)	Stream miles regulated			Wetland acres regulated		
			<i>Rapanos</i> (%) (4)	NWPR (%) (5)	Difference NWPR - <i>Rapanos</i> (6)	<i>Rapanos</i> (%) (7)	NWPR (%) (8)	Difference NWPR - <i>Rapanos</i> (9)
All National	3,154,480	119,825,268	—	—	-686,257	—	—	-34,659,988
Alabama	72,650	4,043,348	0.89	0.87	-1,443	0.86	0.63	-915,215
Arizona	139,281	262,281	0.10	0.01	-12,562	0.33	0.06	-69,684
Arkansas	78,496	2,558,428	0.89	0.67	-16,885	0.77	0.39	-978,770
California	173,028	2,789,804	0.58	0.13	-78,689	0.68	0.15	-1,460,431
Colorado	93,255	1,522,952	0.45	0.17	-26,541	0.43	0.18	-387,822
Connecticut	5,215	304,750	1.00	0.99	-40	0.98	0.48	-152,838
Delaware	2,234	290,940	1.00	0.95	-120	0.99	0.78	-63,450
Florida	22,385	12,681,770	0.99	0.83	-3,636	0.88	0.24	-8,196,536
Georgia	64,833	6,396,737	0.95	0.92	-1,791	0.91	0.47	-2,772,997
Idaho	94,753	1,119,249	0.66	0.40	-24,510	0.68	0.35	-370,204
Illinois	67,074	1,271,986	0.68	0.66	-1,746	0.54	0.42	-147,421
Indiana	24,066	1,008,100	0.72	0.73	286	0.29	0.25	-37,603
Iowa	67,717	1,014,174	0.82	0.71	-7,654	0.63	0.34	-297,352
Kansas	118,236	1,349,856	0.87	0.42	-53,764	0.79	0.33	-620,966
Kentucky	45,616	430,781	0.20	0.30	4,421	0.38	0.29	-35,465
Louisiana	43,096	8,092,819	0.87	0.66	-8,932	0.80	0.55	-2,055,459
Maine	24,974	2,569,961	0.99	0.72	-6,931	0.83	0.32	-1,317,269
Maryland	10,263	863,198	0.98	0.92	-589	0.98	0.66	-278,264
Massachusetts	7,273	775,106	0.99	0.92	-477	0.96	0.41	-425,813
Michigan	47,861	7,712,081	0.55	0.29	-12,435	0.06	0.03	-214,072
Minnesota	60,103	9,973,334	0.11	0.11	-40	0.01	0.01	-58,149
Mississippi	77,386	4,534,181	0.65	0.55	-7,445	0.68	0.43	-1,092,420
Missouri	95,347	1,388,966	0.82	0.63	-17,551	0.78	0.43	-481,217
Montana	166,847	1,589,844	0.77	0.43	-55,447	0.46	0.24	-344,039
Nebraska	72,506	549,755	0.61	0.39	-16,093	0.27	0.16	-58,510
Nevada	143,616	1,003,174	0.23	0.06	-25,156	0.62	0.27	-353,148
New Hampshire	9,374	384,706	0.87	0.54	-3,090	0.75	0.20	-214,531
New Jersey	7,128	1,019,092	1.00	0.99	-81	0.99	0.83	-158,687
New Mexico	109,260	383,873	0.14	0.02	-12,810	0.23	0.05	-68,753
New York	48,756	2,651,158	0.95	0.81	-6,868	0.70	0.30	-1,079,833
North Carolina	56,673	4,679,517	1.00	0.97	-1,535	0.99	0.63	-1,688,348
North Dakota	59,514	2,442,160	0.96	0.78	-10,620	0.20	0.15	-122,294
Ohio	54,736	715,219	0.62	0.68	2,796	0.38	0.26	-88,107

Continued next page

State (1)	Total Stream miles (2)	Total Wetland Acres (3)	Stream miles regulated			Wetland acres regulated		
			<i>Rapanos</i> (%) (4)	NWPR (%) (5)	Difference NWPR - <i>Rapanos</i> (6)	<i>Rapanos</i> (%) (7)	NWPR (%) (8)	Difference NWPR - <i>Rapanos</i> (9)
Oklahoma	75,615	1,274,713	0.89	0.37	-39,658	0.89	0.47	-538,402
Oregon	102,984	1,803,096	0.80	0.43	-37,975	0.88	0.37	-915,645
Pennsylvania	51,477	588,835	0.91	0.77	-7,319	0.78	0.37	-239,928
Rhode Island	978	86,061	1.00	0.94	-61	0.99	0.33	-56,368
South Carolina	29,372	4,238,935	1.00	0.96	-1,019	0.98	0.55	-1,823,810
South Dakota	96,965	3,529,693	0.83	0.57	-24,992	0.35	0.22	-482,699
Tennessee	59,244	1,148,777	0.52	0.46	-3,425	0.56	0.47	-101,837
Texas	176,194	5,551,483	0.67	0.21	-82,033	0.72	0.35	-2,051,586
Utah	82,724	624,397	0.34	0.08	-21,435	0.41	0.14	-169,600
Vermont	7,100	287,628	0.89	0.46	-3,048	0.60	0.15	-128,195
Virginia	49,280	1,682,396	1.00	0.98	-715	0.99	0.86	-219,712
Washington	68,964	1,297,395	0.98	0.60	-26,192	0.97	0.52	-588,074
West Virginia	30,572	81,858	0.50	0.66	4,900	0.65	0.49	-13,097
Wisconsin	53,370	7,610,528	0.59	0.47	-6,227	0.10	0.06	-251,016
Wyoming	106,082	1,646,169	0.49	0.21	-29,080	0.45	0.16	-474,353

Table S10: Deregulation of Drinking Water Areas after NWPR. A 12-digit hydrologic unit code (HUC12) or subwatershed is the finest polygon delineation of watershed boundaries the US Geological Survey defines, corresponding to about 80,000 HUC12s. This table considers active 2019 community water systems (CWS). Column (1) shows the number of people with a CWS served by a HUC12 with at least one point of a given classification deregulated between *Rapanos* and NWPR, divided by the number of people in a CWS served by a HUC12 with at least one point. Column (2) counts number of systems rather than number of people. Column (3) shows the share of points of a given classification that are deregulated in the sample of HUC12's that serve as a drinking water intake. With points as an analogue for water resource miles, 30% of NHD or NWI points deregulated in drinking water systems indicates a deregulation of 30% of NHD and NWI resources that serve as a drinking water source. Column (4) shows the share of points weighted by population. Including people or systems downstream of any HUC12 with at least one deregulated point does not change qualitative results.

Deregulated share of	Population	PWSs	Points serving PWSs	Population Weighted Points
	(1)	(2)	(3)	(4)
A. Same subwatershed (HUC12).				
Deregulated points	0.91	0.85	0.20	0.26
Deregulated NHD or NWI points	0.80	0.69	0.30	0.32
Deregulated NHD points	0.32	0.20	0.21	0.19
Deregulated NWI points	0.80	0.68	0.30	0.32
Deregulated NWI wetlands points	0.55	0.43	0.32	0.46

References and Notes

1. S. M. P. Sullivan, M. C. Rains, A. D. Rodewald, W. W. Buzbee, A. D. Rosemond, Distorting science, putting water at risk. *Science* **369**, 766–768 (2020). [doi:10.1126/science.abb6899](https://doi.org/10.1126/science.abb6899)
[Medline](#)
2. US Environmental Protection Agency and US Army Corps of Engineers, “Economic Analysis for the Final “Revised Definition of `Waters of the United States”” Rule” (2022); https://www.epa.gov/system/files/documents/2022-12/2022_WOTUS%20EA_FinalRule_508.pdf.
3. C. Taylor, H. Druckenmiller, Wetlands, Flooding, and the Clean Water Act. *Am. Econ. Rev.* **112**, 1334–1363 (2023). [doi:10.1257/aer.20210497](https://doi.org/10.1257/aer.20210497)
4. S. Zellmer, Treading water while congress ignores the nation’s environment. *Notre Dame Law Rev.* **88**, 2323 (2013); <https://scholarship.law.nd.edu/ndlr/vol88/iss5/12/>.
5. C. Davenport, “Trump Removes Pollution Controls on Streams and Wetlands,” *New York Times*, 22 January 2020.
6. S. McFetridge, “Judge temporarily blocks clean water rule in 24 states,” AP News, 12 April 2023.
7. US Environmental Protection Agency and US Army Corps of Engineers, “Mapping and the Navigable Waters Protection Rule” (2020); https://19january2021snapshot.epa.gov/sites/static/files/2020-01/documents/nwpr_fact_sheet_-_mapping.pdf.
8. A. Wittenberg, K. Bogardus, “EPA claims ‘no data’ on impact of weakening water rule. But the numbers exist” (E&E News, 2018); <https://www.science.org/content/article/epa-claims-no-data-impact-weakening-water-rule-numbers-exist>.
9. K. A. Fesenmyer, S. J. Wenger, D. S. Leigh, H. M. Neville, Large portion of USA streams lose protection with new interpretation of Clean Water Act. *Freshw. Sci.* **40**, 252–258 (2021). [doi:10.1086/713084](https://doi.org/10.1086/713084)
10. *Sackett v. EPA*. “Brief *amicus curiae* of the National Association of Wetland Managers, the Association of Floodplain Managers, the American Planning Association, the American Water Works Association, and the New England Interstate Water Pollution Control Commission” (2022); https://www.supremecourt.gov/DocketPDF/21/21-454/228256/20220617104921002_21-454%20Brief.pdf.
11. US Army Corps of Engineers, “Compliance with the Federal Endangered Species Act” (2023); <https://www.spk.usace.army.mil/Missions/Regulatory/Permitting/Endangered-Species-Agency-Consultations/>.
12. K. He, X. Zhang, S. Ren, J. Sun, “Deep Residual Learning for Image Recognition” in *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, Las Vegas, NV, USA (2016), pp. 770–778; <https://doi.org/10.1109/CVPR.2016.90>.
13. Materials and methods are available as supplementary materials.
14. L. M. Cowardin, V. Carter, F. C. Golet, E. T. LaRoe, “Classification of wetlands and deepwater habitats of the United States” (FWS/OBS-79/31, U.S. Fish and Wildlife Service, 1979).

15. US Department of Agriculture, National Agriculture Imagery Program (January 2023); https://datagateway.nrcs.usda.gov/GDGHome_DirectDownload.aspx.
16. US Department of Agriculture, Gridded National Survey Geographic Database (October 2022); <https://www.nrcs.usda.gov/resources/data-and-reports/gridded-national-soil-survey-geographic-database-gnatsgo>.
17. US Environmental Protection Agency, National Hydrography Dataset Plus V2 (October 2022); <https://www.epa.gov/waterdata/nhdplus-national-data>.
18. S. A. R. Colvin, S. M. P. Sullivan, P. D. Shirey, R. W. Colvin, K. O. Winemiller, R. M. Hughes, K. D. Fausch, D. M. Infante, J. D. Olden, K. R. Bestgen, R. J. Danehy, L. Eby, Headwater Streams and Wetlands are Critical for Sustaining Fish, Fisheries, and Ecosystem Services. *Fisheries* **44**, 73–91 (2019). [doi:10.1002/fsh.10229](https://doi.org/10.1002/fsh.10229)
19. US Fish & Wildlife Service, National Wetlands Inventory (October 2022); <https://www.fws.gov/program/national-wetlands-inventory/download-state-wetlands-data>.
20. US Army Corps of Engineers, US Army Corps of Engineers Regulatory Boundary (January 2023); https://geospatial-usace.opendata.arcgis.com/datasets/70805e1a8fd74e42b0a9585088d6d151_0/about.
21. US Army Corps of Engineers, “Corps of Engineers Wetlands Delineation Manual” (Y-87-1, 1987); <https://usace.contentdm.oclc.org/digital/collection/p266001coll1/id/4530>.
22. A. Altmann, L. Toloşi, O. Sander, T. Lengauer, Permutation importance: A corrected feature importance measure. *Bioinformatics* **26**, 1340–1347 (2010). [doi:10.1093/bioinformatics/btq134](https://doi.org/10.1093/bioinformatics/btq134) [Medline](#)
23. USCS § 33 1804 (2023).
24. US Environmental Protection Agency, “Land-Use Scenarios: National-Scale Housing-Density Scenarios Consistent with Climate Change Storylines” (EPA/600/R-08/076F, 2009); <https://assessments.epa.gov/iclus/document/&deid=203458>.
25. T. E. Dahl, “Status and Trends of Wetlands in the Conterminous United States 2004 to 2009” (U.S. Fish & Wildlife Service, 2011); <https://www.fws.gov/wetlands/documents/Status-and-Trends-of-Wetlands-in-the-Conterminous-United-States-2004-to-2009.pdf>.
26. J. Kleinberg, H. Lakkaraju, J. Leskovec, J. Ludwig, S. Mullainathan, Human Decisions and Machine Predictions. *Q. J. Econ.* **133**, 237–293 (2018). [Medline](#)
27. J. Podgorski, M. Berg, Global threat of arsenic in groundwater. *Science* **368**, 845–850 (2020). [doi:10.1126/science.aba1510](https://doi.org/10.1126/science.aba1510) [Medline](#)
28. D. M. Katz, M. J. Bommarito 2nd, J. Blackman, A general approach for predicting the behavior of the Supreme Court of the United States. *PLOS ONE* **12**, e0174698 (2017). [doi:10.1371/journal.pone.0174698](https://doi.org/10.1371/journal.pone.0174698) [Medline](#)
29. S. Greenhill, H. Druckenmiller, S. Wang, D. A. Keiser, M. Giroto, J. K. Moore, N. Yamaguchi, A. Todeschini, J. S. Shapiro, (2023). Code for Machine Learning Predicts Which Rivers, Streams, and Wetlands the Clean Water Act Regulates [Dataset], Zenodo; <https://doi.org/10.5281/zenodo.10108709>.

30. US Geological Survey, 3-Dimensional Elevation Program (November 2022); <https://www.usgs.gov/3d-elevation-program>.
31. US Environmental Protection Agency, U.S. EPA Ecoregions, Level IV Ecoregion (January 2023); <https://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>.
32. Prism Climate Group, Parameter-elevation Regressions on Independent Slopes Model, 30-year Normals (January 2023); <https://prism.oregonstate.edu/normals/>.
33. US Geological Survey, National Land Cover Database (December 2022); <https://www.usgs.gov/centers/eros/science/national-land-cover-database#overview>.
34. US Census Bureau, TIGER/Line Shapefiles (December 2022); <https://www2.census.gov/geo/tiger/TIGER2022/>.
35. S. Greenhill, H. Druckenmiller, S. Wang, D. A. Keiser, M. Giroto, J. K. Moore, N. Yamaguchi, A. Todeschini, J. S. Shapiro, (2023). Training data from: Machine Learning Predicts Which Rivers, Streams, and Wetlands the Clean Water Act Regulates [Dataset], Dryad; <https://doi.org/10.5061/dryad.m63xsj47s>.
36. S. Greenhill, H. Druckenmiller, S. Wang, D. A. Keiser, M. Giroto, J. K. Moore, N. Yamaguchi, A. Todeschini, J. S. Shapiro, Prediction data from: Machine Learning Predicts Which Rivers, Streams, and Wetlands the Clean Water Act Regulates [Dataset], Dryad (2023); <https://doi.org/10.5061/dryad.z34tmpgm7>.
37. E. Rolf, J. Proctor, T. Carleton, I. Bolliger, V. Shankar, M. Ishihara, B. Recht, S. Hsiang, A generalizable and accessible approach to machine learning with global satellite imagery. *Nat. Commun.* **12**, 4392 (2021). [doi:10.1038/s41467-021-24638-z](https://doi.org/10.1038/s41467-021-24638-z) [Medline](#)
38. US Environmental Protection Agency, “Clean Water Act Approved Jurisdictional Determinations” (2022); <https://watersgeo.epa.gov/cwa/CWA-JDs/>.
39. U.S. Environmental Protection Agency and Department of the Army, “Economic Analysis for the Proposed “Revised Definition of ‘Waters of the United States’ Rule” (2021); https://www.epa.gov/system/files/documents/2021-11/revised-definition-of-wotus_nprm_economic-analysis.pdf.
40. USDA, “NAIP: Options and Challenges” (2007); https://www.fsa.usda.gov/Internet/FSA_File/2007_esri_uc_naip.pdf.
41. C. Nolte, High-resolution land value maps reveal underestimation of conservation costs in the United States. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 29577–29583 (2020). [doi:10.1073/pnas.2012865117](https://doi.org/10.1073/pnas.2012865117) [Medline](#)
42. US Army Corps of Engineers, “Minimum Standards for Acceptance of Preliminary Wetlands Determinations” (2001); <https://www.spk.usace.army.mil/Portals/12/documents/regulatory/pdf/Minimum%20Standards.pdf>.
43. US Army Corps of Engineers, “Jurisdictional Determinations and Delineating Waters of the United States, including Wetlands” (2013); https://www.spk.usace.army.mil/Portals/12/documents/regulatory/Reg_workshop/RPW_JDs_and_Delineation.pdf.

44. Federal Register, “Revised Definition of ‘Waters of the United States’” (EPA-HQ-OW-2021-0602, 2023); <https://www.federalregister.gov/documents/2023/01/18/2022-28595/revised-definition-of-waters-of-the-united-states>.
45. *Sackett v. Environmental Protection Agency*, 598 U.S. ____ (2023); https://www.supremecourt.gov/opinions/22pdf/21-454_4g15.pdf.
46. US Environmental Protection Agency and US Army Corps of Engineers, “Amendments to the ‘Revised Definition of ‘Waters of the United States’” (EPA-6560-50-P, 2023); <https://www.epa.gov/system/files/documents/2023-08/Pre-publication%20Version%20of%20the%20Final%20Rule%20-%20Amendments%20to%20the%20Revised%20Definition%20of%20Waters%20of%20the%20United%20States.pdf>.